

DISCOVERY OF A NUCLEAR X-RAY SOURCE IN NGC 7331: EVIDENCE FOR A MASSIVE BLACK HOLE

Christopher J. Stockdale, W. Romanishin, and John J. Cowan

Department of Physics and Astronomy, University of Oklahoma, Norman, OK 73019

stockdal@mail.nhn.ou.edu,wjr@mail.nhn.ou.edu,cowan@mail.nhn.ou.edu

Received _____; accepted _____

ABSTRACT

Using ROSAT, we have made the first deep X-ray observations of NGC 7331, discovering a nuclear X-ray source coincident with the previously determined radio and optical nuclear centers. Positions, luminosities, and fluxes of X-ray sources in the field are compared to previously identified radio and optical sources. The nucleus of NGC 7331 has been analyzed to discern any new evidence supporting the presence of a massive black hole (MBH), and comparisons are made with other Low-Ionization Nuclear Emission-line Region (LINER) galaxies. The flux ratio of core radio to soft X-ray emission in NGC 7331 is lower than in other LINERs included in our sample.

Subject headings: galaxies: individual (NGC 7331, NGC 7335) — galaxies: nuclei — X-rays: galaxies

1. Introduction

Deep radio observations of NGC 7331 were first made in an attempt to search for the supernova remnant (SNR) from SN1959D (Cowan, Romanishin, & Branch 1994, hereafter = CRB). Although the SNR was not identified in those observations, an unresolved nuclear source was detected (CRB). Optical observations of the nucleus of NGC 7331 identified an optical LINER spectrum (Keel 1983; Bower 1992). These two discoveries support the possibility that NGC 7331 harbors an MBH. Rubin et al. (1965) and Afanasiev, Silchenko, & Zasov (1989) made kinematic studies, measuring rotational motion from [NII] and H α emission lines. Based on these observations were analyzed by Afanasiev et al.(1989) suggested an MBH of $\sim 5 \times 10^8 M_{\odot}$ in NGC 7331. Bower et al. (1993), however, suggested that this rotational motion could be explained by models without an MBH and with a constant mass-to-light ratio as a function of radius. Bower et al. (1993) were only able to exclude an MBH with a mass greater than $5 - 10 \times 10^8 M_{\odot}$ (CRB), a result which is relatively insensitive to small differences in the assumed distance to the galaxy. The Bower et al. (1993) mass limit is also near the upper limit of measured masses for Massive Dark Objects found in LINER galaxies, which have X-ray and radio properties similar to those found in NGC 7331 (Ho 1998).

In this Letter we report on the first deep ROSAT X-ray observations of NGC 7331 and their impact in understanding the nature of its LINER nucleus. In §2, we discuss the results of the observations and the techniques used to reduce these data. Classically, the possible indicators for an MBH are the detection of an unresolved, nuclear source, a non-thermal radio and X-ray spectral index, variability in the nuclear flux over months to years, and an anti-correlation between X-ray and radio variability (Melia 1992). The fact that NGC 7331 possesses an unresolved nuclear source and has a non-thermal radio spectral index was confirmed by CRB. This Letter identifies a similarly unresolved X-ray source, but can make

no statement as to the nature of the X-ray spectra. Confirmation of a non-thermal X-ray spectra and the last two criteria for an MBH require further X-ray and radio observations. While starburst galaxies can also have non-thermal, unresolved sources, there is no evidence for an anti-correlation in X-ray and radio variability in such galaxies. Other indicators of the presence of an MBH in NGC 7331 will be discussed later in §3 of this Letter. Also in §3, we will provide comparisons of our results with current theories and observations of low luminosity AGN (LLAGN) powered LINERs and make our conclusions.

2. Observations and Data Reduction

NGC 7331 is an Sbc spiral galaxy, which was observed with the ROSAT High Resolution Imager (HRI) in December 1995 to study the nuclear source that had been previously observed in the radio with the Very Large Array (VLA) (CRB). The HRI is an X-ray photon collector with no spectral resolution, collecting photons with energies ranging from 0.12 keV to 2.4 keV. The observations were made over a two day period with a total observation time of 30.5 ks. The search was centered on the CRB radio center of NGC 7331 at R.A.(2000) = $22^h37^m4^s8$, Decl.(2000) = $+34^\circ25'11''.5$.

Raw data reduction was provided by the ROSAT Standard Data Processing Center. Standard ROSAT tests for short term variability are not applicable due to insufficient counts. The data were further reduced using the IRAF routine GAUSS which convolved the data with a circularly symmetric Gaussian function, with the parameter sigma set to one pixel, effectively $8''.0$. The image was smoothed with a Gaussian because of concerns from the Data Processing Center that the standard analysis system was not very reliable in properly identifying sources in complex emission regions. Sources that were initially reported as individual point sources by the Center, are actually part of a more complex X-ray emission region. The data were then evaluated with the NOAO-IRAF routine

QPHOT to obtain object counts, count noise, and accurate position measurements. The aperture radius used to determine the count flux from the galaxy was set to $24''$, and sky background measurements were determined with the specified parameters of an inner annulus of $80''$ and outer annulus of $102''$.

Lacking direct X-ray spectral data, a value for the $\log N(H) = 21.3$ (Burstein & Heiles 1978; van Steenberg et al. 1988; Stark et al. 1992; Smith 1998) and a photon index, Γ ($dF/dE \propto E^{-\Gamma}$), of 1.0 and 2.0 for the central sources in NGC 7331 were assumed to determine the unabsorbed X-ray flux. The total column density of hydrogen includes both the Galactic column density (Burstein & Heiles 1978) and the column density for the bulge of NGC 7331 (Smith 1998). The ROSAT HRI energy-to-count conversion factors were taken from the ROSAT Users Handbook, assuming a power law spectrum. The assumption of an X-ray power law spectrum is consistent with both the possible presence of an MBH, as well as for a post starburst galaxy (Fabbiano 1996). The X-ray luminosity for the galaxy was calculated assuming the distance to NGC 7331 as 15.1 Mpc (Hughes et al. 1998). The detections reported in Table 1 all meet at least a 3σ detection threshold. Since little is known about the other detected sources in Table 1, a power law spectrum, a photon index of 1.0 and the same line-of-sight, Galactic column density, which was used for NGC 7331, were used to determine their energy flux measurements.

Sources of uncertainty in ROSAT positions include a known attitude solution error, which causes X-ray position offsets of order $6''$ (ROSAT Users Guide). Since any correction would require knowing X-ray positions and this is the first high resolution image of the field, it is not possible to accurately correct for this error. Table 3 compares the optical, radio, and X-ray positions of the nucleus of NGC 7331. The values for the nuclear positions agree to well within $5''$ and are definitely coincident within the acceptable ROSAT error.

3. Results and Discussion

The observations of the X-ray nuclear source in NGC 7331 support the likelihood of an MBH in its core. In Table 2, we compare the luminosity of the nuclear source in NGC 7331 with luminosity values for other LINER galaxies. Also included in this table for comparison are the reported luminosities for two, identified starburst galaxies.

An assumed power-law spectrum and photon indices of 1.0 and 2.0 were used to analyze our ROSAT observations of NGC 7331, because these are typical of other observed X-ray LINERs in the energy range from 0.1 keV to 2.4 keV. The X-ray luminosity of NGC 7331 in this bandpass region is within the observed range found in other LINERs and somewhat larger than those luminosities observed in typical starburst galaxies. We note that these LINER X-ray luminosities are near the upper limit of X-ray luminosities typically found in normal spiral galaxies (Fabbiano 1996), and therefore an X-ray luminosity cannot be used solely to identify possible MBHs.

The radio spectrum of a typical LLAGN is predicted to be powered by cyclotron synchrotron emissions in the advection dominated accretion flow (ADAF) model used by Mahadevan (1997). The observed spectral index, α , of -0.6 ($S_\nu \propto \nu^{+\alpha}$) for NGC 7331 indicates a non-thermal source powering the radio spectrum between 6 cm and 20 cm and is in agreement with observations of similar sources listed in Table 4. The value reported by CRB for the spectral index of NGC 7331 agrees closely with the theoretical value of the spectral index of M31, which was modeled with an MBH by Melia (1992). It should also be noted that this value of α is only slightly greater than the spectral index normally associated with SNRs, -0.8 , and this might suggest a post starburst nature (Condon 1996).

Table 4 also lists radio fluxes from other LINERs with identified nuclear radio sources. The same value of the spectral index found for NGC 7331 is reported for M51 and M81 by Turner & Ho (1994), and M81 has been confirmed to harbor an LLAGN by Ho et

al. (1996). While the CRB spectral index alone does not confirm the exact nature of the nuclear source in NGC 7331, it does suggest that NGC 7331 is an LLAGN, given the galaxy’s identification as a LINER galaxy and the similarity of its spectral index with other LINER galaxies confirmed to harbor MBHs.

For a sample of LINER nuclei Table 5 lists the ratio of soft X-ray and 6 cm radio fluxes. The X-ray fluxes listed in Table 5 are averaged over their stated bandpasses listed in Table 2. Among this limited sample, there appears to be a range for galaxies, from being relatively radio quiet (or X-ray loud) ($\log [\text{X-Ray}/\text{Radio}] \propto -1.8$) to relatively radio loud (or X-ray quiet) ($\log [\text{X-Ray}/\text{Radio}] \propto -4.6$). It must be noted that this is a small sample and that these observations were taken at various epochs and with different instruments. (Radio observations are from the VLA and the Westerbork Synthesis Radio Telescope and X-ray observations are from ROSAT and ASCA.)

Another predicted indicator of an MBH is the bipolar outflow, which is believed to develop as the result of convective instabilities in the thin disk approximation. This in turn leads to a quasi-spherical accretion flow, characteristic of the advective model (Narayan & Yi 1995). In the case of NGC 7331, these outflows have not been positively identified in any observations to date. If these features do exist, they are likely to extend only a few parsecs. For NGC 3079, a LINER galaxy 16 Mpc away, jets were identified which extended only 1.5 pc from the central engine (Trotter et al. 1998). Trotter et al. (1998) detected a molecular disk with a binding mass of $\sim 10^6 M_\odot$. The radio observations of NGC 7331 indicate only a compact nuclear source with a ring of non-thermal sources extending beyond the optical galaxy (CRB). The fact that similar jets have not been detected in NGC 7331 can be attributed to the galaxy being 15.1 Mpc away ($1'' = 73$ pc), which makes resolution of parsec-scale outflows beyond the capability of the CRB observations.

Further evidence supporting the existence of a LINER nucleus in NGC 7331, similar to

the one in M31, has been presented by Mediavilla et al. (1997) and further confirmed by Heckman (1996). M31 was identified as harboring an MBH of $3 \times 10^7 M_{\odot}$ by Kormendy & Richstone (1995). Mediavilla et al. (1997) and Ciardullo et al. (1988) report that the kinematics of the stars and ionized gas are decoupled, in NGC 7331 and M31 respectively. This is in contrast to the conditions found in Seyfert galaxies. Also, Tosaki & Shioya (1997) and Young & Scoville (1982) found no CO emission near the nuclei of NGC 7331 and M31. This is atypical for post starburst galaxies. Their results do support the possibility that this absence of CO emission may be due to the presence of an MBH.

Also identified in the HRI image was a source which is within $9''$ of the optical position of NGC 7335 (Klemola et al. 1987). The observed flux was $2.15 \times 10^{-13} \text{ erg sec}^{-1} \text{ cm}^{-2}$ (assuming a power-law spectrum, a photon index of 1.0, and the same line of sight, Galactic column density of hydrogen used in the previous section). If this X-ray source is associated with NGC 7335, this would indicate an X-ray luminosity of $3.1 \times 10^{41} \text{ erg s}^{-1}$, assuming a distance to NGC 7335 of 110 Mpc ($z=6315 \text{ km/s}$ de Vaucouleurs et al. (1991) and a Hubble Constant of $55 \text{ km s}^{-1} \text{ Mpc}^{-1}$). Few observations have been made of this galaxy, and little is known about it. Based upon this derived X-ray luminosity, however, this galaxy warrants further study to determine if it is also a possible candidate for AGN activity.

There are a number of tests associated with X-ray studies that could more definitively determine the nature of the nucleus of NGC 7331 (see e.g., Fabbiano 1996; Serlemitsos et al. 1996; Nandra et al. 1997; Ho 1998). Such tests will require deep observations, which could provide resolution of the spectrum at multiple wavelengths and identify any form of variability in the observed flux from NGC 7331.

The research was supported in part by NASA Grant NAG5-3214 and has made use of the NASA/IPAC Extragalactic Database (NED), which is operated by the Jet Propulsion Laboratory, Caltech, under contract with the National Aeronautics and Space

Administration. We thank an anonymous referee for helping us to improve this Letter.

REFERENCES

- Afanasiev, V. I., Silchenko, O. K., & Zasov, A. V. 1989, *A&A*, 213, L9
- Argyle, R. W. & Clements E. D. 1990, *Observatory*, 110, 93
- Bower, G. A. 1992, PhD Thesis, Univ. Michigan
- Bower, G. A., Richstone, D. O., Bothum, G. D., & Heckman, T. M. 1993, *ApJ*, 402, 76
- Burstein, D. & Heiles C. 1984, *ApJS*, 54, 33
- Ciardullo, R., Rubin, V., Jacoby, G. H., Ford, H. C., & Ford W. K. 1988, *AJ* 95, 438
- Condon, J. J. 1996, *ASP Conf*, 103, 132.
- Cowan, J.J., Romanishin, W., & Branch D. 1994, *ApJ*, 436, L139 (CRB)
- Crane, P. C., Cowan, J. J., Dickel, J. R., & Roberts, D. A. 1993, *ApJ*, 417, L61
- Crane, P. C., Dickel, J. R., & Cowan, J. J. 1992, *ApJ*, 390, L9
- Cui, W., Feldkhun, D., & Braun, R. 1997, *ApJ*, 477, 693
- Dahlem, M., Heckman, T. M., & Fabbiano, G. 1995, *ApJ*, 442, L49
- de Vaucouleurs, G., de Vaucouleurs, A, Corwin, H. G., Buta, R. J., Paturel, G., Fouqué, P.
1991, *Third Reference Catalog of Bright Galaxies* Springer-Verlag, New York (RC3)
- Fabbiano, G. 1996, *ASP Conf*, 103, 56
- Heckman, T. 1996, *ASP Conf*, 103, 241
- Ho, L. C., Filippenko, A. V., & Sargent W. L. W. 1996, *ApJ* 462, 18.
- Ho, L. C. 1998, *Observational Evidence for Black Holes in the Universe*, in print

- Hughes, S. M. G., Han, M., Hoessel, J., Freeman, W. L., Kennicutt, R. C., Mould, J. R., Saha, A., Stetson, P. B., Madore, B. F., Silbermann, N. A., Harding, P., Ferrarese, L., Ford, H., Gibson, B. K., Graham, J. A., Hill, R., Huchra, J., Illingworth, G. D., Phelps, R., & Sakai, S. 1998 *ApJ*, 501, 32
- Hummel, E. 1980, *A&AS*, 41, 151
- Keel, W. C. 1983, *ApJ*, 269, 466
- Klemola, A. R., Jones, B. F., & Hanson, R. B., 1987, *AJ*, 94, 501
- Koratkar, A., Deustus. S. E., Heckman, T., Filippenko, A. V., Ho, L. C., & Rao, M., 1995, *ApJ*, 440, 132
- Kormendy, J. & Richstone, D. 1995, *ARA&A*, 33, 581
- Long, K. S., Charles, P. A., Blair, W. P., & Gordon, S. M., 1996, *ApJ*, 466, 750
- Mahadevan, R. 1997, *ApJ*, 477, 585
- Mediavilla, E., Arribas, S., García-Lorenzo, B., & del Brugo, C. 1997, *ApJ*, 488, 682
- Melia. F. 1992, *ApJ*, 398, L95
- Nandra, K., George, I. M., Mushotzky, R. F., Turner, T. J., & Yaqoob, T. 1997, *ApJ*, 477, 602
- Narayan, R. & Yi, I. 1995, *ApJ*, 444, 231
- Pietsch, W., Volger, A., Jain, A., & Klein, U. 1994, *A&A*, 284, 386
- ROSAT Users Guide, Max-Planck-Institut, 1994
- Rubin, V. C., Burbidge, E. M., Burbidge, G. R., & Crampin, D. J. 1965, *ApJ*, 141, 759

- Serlemitsos, P., Ptak, A., & Yaqoob, T. 1996, ASP Conf., 103, 70
- Smith, B. J. 1998, ApJ, 500, 181
- Stark, A. A., Gammie, C. F., Wilson, R. W., Bally, J., Linke, R. A., Heiles, C., & Hurwitz, M. 1992, ApJS, 79, 77
- Tosaki, T. & Shioya, Y. 1997, ApJ, 484, 664
- Trotter, A. S., Greenhill, L., J., Moran, J. M., Reid, M. J., Irwin, J. A. & Lo, K. 1998, ApJ, 495, 740
- Turner, J. L., & Ho, P. T. P. 1994, ApJ, 421, 122
- van Steenberg, M. E. & Shull, J. M. 1988, A Decade of UV Astronomy with the IUE Satellite, Vol. 2 (Noordwijk, The Netherlands: ESTEC), 219
- Vila, M. B., Pedlar, A., Davies, R. D., Hummel, E., & Axon, D. J. 1990, MNRAS, 242, 379
- Young, J. S., & Scoville, N. 1982, ApJ, 260, L41

TABLE 1. X-Ray Sources in the Field of NGC 7331

Source	RA	Decl.	Counts	Flux	Comments
	(2000)	(2000)	(ksec ⁻¹)	(10 ⁻¹³ erg s ⁻¹ cm ⁻²)	
1	22 ^h 37 ^m 3 ^s .64	+34°24′59″.27	4.17 ± .4	41.3 ± .63	NGC 7331
2	22 ^h 37 ^m 1 ^s .86	+34°26′53″.90	1.3 ± .3	2.15 ± .36	NGC 7335
3	22 ^h 36 ^m 29 ^s .38	+34°25′56″.78	0.9 ± .3	1.49 ± .50	
4	22 ^h 36 ^m 33 ^s .13	+34°30′11″.49	1.9 ± .4	3.16 ± .66	
5	22 ^h 37 ^m 28 ^s .97	+34°25′25″.49	1.0 ± .3	1.66 ± .50	
6	22 ^h 37 ^m 38 ^s .15	+34°22′27″.36	0.5 ± .3	0.83 ± .50	
7	22 ^h 37 ^m 3 ^s .48	+34°9′47″.85	1.4 ± .7	2.33 ± 1.16	
8	22 ^h 37 ^m 51 ^s .93	+34°28′29″.43	0.9 ± .4	1.49 ± .66	
9	22 ^h 35 ^m 54 ^s .60	+34°13′51″.34	9.2 ± .9	15.3 ± 1.5	
10	22 ^h 36 ^m 3 ^s .43	+34°12′14″.95	5.9 ± .9	9.80 ± 1.50	

TABLE 2. Comparisons of NGC 7331 and
Other Similar X-ray Sources

Source	Type	Luminosity	Photon Index	Bandpass	Mechanism	Ref
	(RC3)	$10^{40} \text{ erg s}^{-1}$		keV		
NGC 7331	SAab	2.70 — 8.46	$1.0 < \Gamma < 2.0$	0.1-2.4	LINER	this Letter
NGC 4258	SABbc	0.51	3.7	0.1-2.4	LINER	1
NGC 4736	R SA(r)ab	0.34	2.3	0.1-2.0	LINER	2
NGC 4594	SAa	5.3	1.80	0.5-2.0	LINER	3
M51	SABc	2.5	1.76	0.5-2.0	LINER	3
NGC 3079	SBc	4.4	1.76	0.5-2.0	LINER	3
M81	SAab	1.2	1.9	0.5-2.0	LINER	3
NGC 3642	SA(r)bc	3.9	2.84	0.2-2.2	LINER	4
NGC 3147	SA(r)bc	24	1.74	0.5-2.0	LINER	3
NGC 4579	SAB(rs)b	28	1.87	0.5-2.0	LINER	3
M33	SACd	0.1248	0.74	0.1-2.4	Starburst	5
NGC 3628	SAb	< 0.06 —1.7	0.27—1.16	0.1-2.0	Starburst	6

References

1. Pietsch et al. (1994)
2. Cui et al. (1997)
3. Serlemitsos et al. (1996)
4. Koratkar et al. (1995)
5. Long et al. (1996)
6. Dahlem et al. (1995)

TABLE 3. Nuclear Positions for NGC 7331

Observation	R.A. (2000)	Decl.(2000)	Reference
X-ray	$22^h37^m3^s.64$	$+34^\circ24'59''.27$	this Letter
Radio	$22^h37^m4^s.00$	$+34^\circ24'57''.07$	CRB (1994)
Optical	$22^h37^m4^s.01$	$+34^\circ24'56''.07$	Klemola 1994, private communication
Optical	$22^h37^m4^s.04$	$+34^\circ24'56''.47$	Argyle & Clements (1990)

TABLE 4. Core Flux Comparisons of NGC 7331 and
Other Similar Radio Sources

Source	2cm	3.6cm	6cm	20cm	Spectral Index	Ref
Type	mJy	μ Jy	mJy	mJy		
NGC 7331	...	...	0.121	0.234	−0.6	1
SAab	...	...	(B array)	(A array)		
NCG 4258	3.2	...	2.4	< 10	+0.4	5 & 2
	...	...	1.5	2.0	−0.2	3
SABbc	(C array)	...	(B array)	(see ref) & (A array)		
NCG 4579	...	...	...	40	...	2
SAB(rs)b	...	...	...	(see ref)		
NCG 4736	3.2	...	6.1	27	−0.3	2 & 5
	2.2	...	3.6	...	−0.4	5
R SA(r)ab	(C array)	...	(B array)	(see ref)		
M31	...	28 & 39	...	...	...	4
SA(s)b	...	(A array)	...	...		
M51	2.3	...	1.6	...	−0.6	5
SAbc	(C array)	...	(B array)	...		
M81	45	...	92	...	−0.6	5
SAbc	(C array)	...	(B array)	...		

References

1. CRB
2. Hummel (1980) (Westerbork Synthesis Radio Telescope)
3. Vila et al. (1990)
4. Crane et al. (1992); Crane et al. (1993)
5. Turner & Ho (1994)

TABLE 5. Soft X-ray & 6 cm Radio Flux Ratios for Selected LINERs

Source	Radio Flux ^a	X-ray Flux ^a	log [X-Ray/Radio]	Ref
	10 ⁻²⁷	10 ⁻³⁰		
NGC 7331	0.121	1.93 — 5.90	(−1.80) — (−1.31)	this Letter & 1
NGC 4258	2.4	7.4	−2.1	2 & 3
M51	1.6	3.2	−2.7	5 & 6
NGC 4736	3.6	4.1	−2.9	3 & 4
NGC 4736	6.1	4.1	−3.2	3 & 4
NGC 4579	40	5.3	−3.9	3 & 6
M81	92	2.3	−4.6	5 & 6

a. $\text{erg s}^{-1} \text{ cm}^{-2} \text{ Hz}^{-1}$

References

1. CRB
2. Pietsch et al. (1994)
3. Hummel (1980)
4. Cui et al. (1997)
5. Turner & Ho (1994)
6. Serlemitsos et al. (1996)

Fig. 1.— FIG. 1 A grey scale image of NGC 7331 and the field. NGC 7331 is the circled object labeled #1 and NGC 7335 is the circled object labeled #2. The X-ray detection of the nuclear source in NGC 7331 is within $5''.0$ of prior radio observations of the nucleus given in Table 3. The remaining eight unidentified sources listed in Table 1 (#3 - #10) are indicated on this figure.

